

(19)  Canadian  
Intellectual Property  
Office

An Agency of  
Industry Canada

Office de la Propriété  
Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

(11) **CA 2 463 690** (13) **A1**

(40) 24.04.2003  
(43) 24.04.2003

(12)

(21) 2 463 690

(22) 16.10.2002

(51) Int. Cl.<sup>7</sup>: **A61K 9/16, A61K 47/02,**  
**A61P 1/04, A61K 9/10,**  
**A61K 9/26, A61K 47/26,**  
**A61K 47/32, A61K 47/36,**  
**A61K 47/38, A61K 31/4439,**  
**A61K 9/48, A61K 9/50**

(85) 15.04.2004

(86) PCT/JP02/010720

(87) WO03/032953

(30) 2001-319444 JP 17.10.2001

(71) **TAKEDA CHEMICAL INDUSTRIES, LTD.,**  
1-1 Doshomachi 4-chome  
Chuo-ku, Osaka-shi  
541-0045, OSAKA, XX (JP).

(72) **SHIMIZU, TOSHIHIRO (JP).**  
**NAKANO, YOSHINORI (JP).**

(74) **FETHERSTONHAUGH & CO.**

(54) **GRANULES CONTENANT UN AGENT CHIMIQUE INSTABLE EN MILIEU ACIDE, EN CONCENTRATION ELEVÉE**

(54) **GRANULES CONTAINING ACID-UNSTABLE CHEMICAL IN LARGE AMOUNT**

(57)

It is intended to provide preparations such as capsules containing an acid-unstable chemical (in particular, a benzimidazole compound having an antilulcer effect, etc.) at a high concentration which are prepared by using about 12% by weight or more (based on the total granules) of the acid-unstable chemical and blending a basic inorganic salt therewith to give granules of about 600 .mu.m or more in the average grain size.



Office de la Propriété  
Intellectuelle  
du Canada  
Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office  
An agency of  
Industry Canada

CA 2463690 A1 2003/04/24

(21) **2 463 690**

(12) **DEMANDE DE BREVET CANADIEN  
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2002/10/16

(87) Date publication PCT/PCT Publication Date: 2003/04/24

(85) Entrée phase nationale/National Entry: 2004/04/15

(86) N° demande PCT/PCT Application No.: JP 2002/010720

(87) N° publication PCT/PCT Publication No.: 2003/032953

(30) Priorité/Priority: 2001/10/17 (2001-319444) JP

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> A61K 9/16, A61K 9/50, A61K 9/48,  
A61K 47/02, A61K 31/4439, A61K 47/38, A61K 47/36,  
A61K 47/32, A61K 47/26, A61K 9/26, A61K 9/10,  
A61P 1/04

(71) Demandeur/Applicant:  
TAKEDA CHEMICAL INDUSTRIES, LTD., JP

(72) Inventeurs/Inventors:  
SHIMIZU, TOSHIHIRO, JP;  
NAKANO, YOSHINORI, JP

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : GRANULES CONTENANT UN AGENT CHIMIQUE INSTABLE EN MILIEU ACIDE, EN CONCENTRATION  
ELEVÉE

(54) Title: GRANULES CONTAINING ACID-UNSTABLE CHEMICAL IN LARGE AMOUNT

(57) Abrégé/Abstract

It is intended to provide preparations such as capsules containing an acid-unstable chemical (in particular, a benzimidazole compound having an antulcer effect, etc.) at a high concentration which are prepared by using about 12% by weight or more (based on the total granules) of the acid-unstable chemical and blending a basic inorganic salt therewith to give granules of about 600  $\mu$ m or more in the average grain size.

Canada

<http://opic.gc.ca> · Ottawa-Hull K1A 0C9 · <http://cipo.gc.ca>

OPIC · CIPD 191

OPIC



CIPO

ABSTRACT

It is intended to provide preparations such as capsules containing an acid-unstable medicament (in particular, a benzimidazole compound having an antiulcer effect, etc.) at a high concentration which are prepared by using about 12% by weight or more (based on the total granules) of the acid-unstable chemical and blending a basic inorganic salt therewith to give granules of about 600  $\mu\text{m}$  or more in the average particle size.

## GRANULES CONTAINING ACID-UNSTABLE CHEMICAL IN LARGE AMOUNT

## TECHNICAL FIELD

The present invention relates to stabilized granules,  
5 which comprises a high content of an acid-unstable  
medicament, in particular, a benzimidazole compound useful  
as an antiulcer agent and a basic inorganic substance.

## BACKGROUND ART

10 Since benzimidazole compounds such as lansoprazole,  
omeprazole and rabeprazole have gastric acid secretion  
inhibitory activity, gastric mucosa protecting activity,  
etc., they are widely used as a peptic ulcer treating agent.

However, these compounds are inferior in stability,  
15 and are unstable to humidity, temperature and light. In  
particular, they are unstable to acids and, when formulated  
into an aqueous solution or a suspension, they become  
extremely unstable with lowering of pH.

In addition, stability in the form of preparations,  
20 i.e., tablets, powders, fine granules, capsules, etc.  
becomes lower than that of the compounds alone because of  
their strong interaction with other components in  
formulated preparations, and color change or degradation is  
observed upon production or storage of the preparations.  
25 In order to stabilize them, JP 62-277322 A discloses

enteric granules, enteric fine granules, etc. obtained by blending a stabilizing agent composed of a basic inorganic salt of magnesium and/or calcium, followed by enteric coating.

5           Meanwhile, it is necessary to apply an enteric coating to a benzimidazole compound because the compound has such properties that it is hardly soluble in water and is unstable to acids. An enteric coating does not dissolve in a stomach containing a relatively larger amount of water,  
10 but dissolves in a small intestine containing a smaller amount of water, whereby a benzimidazole compound is dissolved and absorbed. That is, since a composition containing a benzimidazole compound is required to be rapidly disintegrated in a small intestine, granules which  
15 have a larger surface area, and which are more easily and rapidly disintegrated or dissolved than tablets are considered to be more desired.

          In Examples specifically disclosed in JP 62-277322 A, the content of benzimidazole compound is about 6.3 to 11.5%  
20 based on the total enteric granules, and the capsule in which the granules are filled is No. 1 or 2. Pharmacy Vol. 50(3) 230-238 (1990) reports that smaller capsules are more easily administered in view of an appearance organoleptic test, and a limit of a capsule size which can be easily  
25 administered is No. 3. Then, No. 1 or 2 capsule may reduce

compliance of patients, in particular, elderly patients who have difficulty in swallowing.

#### OBJECTS OF THE INVENTION

5           In order to facilitate administration to patients, in particular, elderly or pediatric patients who have difficulty in swallowing, and to improve compliance, an object of the present invention is to produce stable enteric granules which comprises an acid-unstable  
10       medicament including a benzimidazole compound in a high content and a basic inorganic salt, and to render a size of a capsule in which the enteric granules are filled suitable for easy administration.

#### 15       SUMMARY OF THE INVENTION

          The present inventors have found that, in granules which contain an acid-unstable medicament, in particular, a benzimidazole compound and a basic inorganic salt as a stabilizing agent, and which are coated with an enteric  
20       layers or agents, the acid-unstable medicament can be stabilized even in high concentration and a high content, and can be easily administered to patients with improved compliance by adjusting a blending ratio of the basic inorganic salt to the acid-unstable medicament and an  
25       average particle size properly. The present inventors have

further studied and, as a result, the present invention has been completed.


That is, the present invention provides:

(1) Granules comprising a principal ingredient layer  
5 containing an acid-unstable medicament in an amount of about 12% by weight or more based on the total granules; an intermediate coating layer which is formed on the principal ingredient layer; and an enteric coating layer which is formed on the intermediate coating layer; wherein said  
10 granules contain a basic inorganic salt, and have an average particle size of about 600  $\mu\text{m}$  or more;

(2) The granules according to the above (1), wherein the basic inorganic salt is a magnesium salt or a calcium salt;

15 (3) The granules according to the above (1), wherein the acid-unstable medicament is a proton pump inhibitor (PPI);

(4) The granules according to the above (3), wherein the PPI is a benzimidazole compound represented by the  
20 formula (I):



wherein ring A is an optionally substituted benzene ring,

R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are the same or different, and represent a hydrogen atom, an optionally substituted alkyl group, an optionally substituted alkoxy group, or an optionally substituted amino group, and Y represents a nitrogen atom or CH, or a salt thereof;

(5) The granules according to the above (3), wherein the PPI is lansoprazole, omeprazole, rabeprazole, pantoprazole, leminoprazole, tenatoprazole (TU-199), or an optically active isomer thereof, or a pharmaceutically acceptable salt thereof;

(6) The granules according to the above (3), wherein the PPI is lansoprazole, or an optically active isomer thereof, or a pharmaceutically acceptable salt thereof;

(7) The granules according to the above (4), wherein the basic inorganic salt is contained in the principal ingredient layer in an amount of about 0.2 part by weight to about 0.6 part by weight based on 1 part by weight of the benzimidazole compound;

(8) The granules according to the above (1) which are granules having a core, wherein the principal ingredient layer is formed on the core comprising at least one material selected from sucrose, starch, lactose and crystalline cellulose;

(9) The granules according to the above (1), wherein the enteric coating layer contains an enteric water-soluble



polymer;

(10) The granules according to the above (9), wherein the enteric water-soluble polymer is a methacrylic acid copolymer;

5 (11) The granules according to the above (1), wherein an average particle size of the granules is about 1000  $\mu\text{m}$  to about 2000  $\mu\text{m}$ ;

(12) The granules according to the above (4), wherein the benzimidazole compound is contained in an amount of  
10 about 12% by weight to about 40% by weight based on the total granules;

(13) A granular preparation, a capsule, a tablet, an effervescent preparation or a suspension comprising the granules according to the above (1);

15 (14) Granules comprising a principal ingredient layer containing a PPI in an amount of about 12% by weight to about 40% by weight based on the total granules, and one or more basic inorganic salts selected from the group consisting of basic salts of a sodium salt, a potassium  
20 salt, an aluminum salt, a magnesium salt and a calcium salt in an amount of about 0.2 part by weight to about 0.6 part by weight based on 1 part by weight of the PPI; an intermediate coating layer which is formed on the principal ingredient layer; and an enteric coating layer which is  
25 formed on the intermediate coating layer, and having an

average particle size of about 1000  $\mu\text{m}$  to about 2000  $\mu\text{m}$ ;

(15) A granular preparation, a capsule or a tablet comprising the granules according to the above (14);

(16) The preparation according to the above (15),  
5 which is a preparation for treating or preventing peptic ulcer, Zollinger-Ellison syndromes, gastritis, reflux esophagitis, symptomatic gastroesophageal reflux disease (symptomatic GERD), NUD (non ulcer dyspepsia), stomach cancer, gastric MALT lymphoma, ulcer resulting from  
10 nonsteroidal antiinflammatory medicaments, or excess stomach acid or ulcer due to post-operation stress, a preparation for eradicating *Helicobacter pylori*, or a preparation for suppressing upper gastrointestinal hemorrhage due to peptic ulcer, acute stress ulcer,  
15 hemorrhagic gastritis or invasion stress;  
(17) Granules comprising a principal ingredient layer containing lansoprazole or an optically active isomer (R isomer) thereof in an amount of about 14% by weight to about 20% by weight based on the total granules, and one or  
20 more basic inorganic salts selected from the group consisting of basic salts of magnesium and calcium in an amount of about 0.2 part by weight to about 0.4 part by weight based on 1 part by weight of lansoprazole or an optically active isomer (R isomer) thereof; an intermediate  
25 coating layer which is formed on the principal ingredient

layer; and an enteric coating layer which is formed on the intermediate coating layer, and having an average particle size of 1000  $\mu\text{m}$  to about 2000  $\mu\text{m}$ ;

(18) The granules according to the above (17), wherein  
5 the basic inorganic salt is magnesium carbonate;

(19) No. 3 to 5 capsules comprising 30 mg of  
lansoprazole per one capsule;

(20) The capsule according to the above (19), wherein  
the granules according to the above (17) are filled;

10 (21) No. 4 to 5 capsules comprising 15 mg of  
lansoprazole per one capsule;

(22) The capsule according to the above (21), wherein  
the granules according to the above (17) are filled;

(23) No. 1 to 3 capsules comprising 60 mg of  
15 lansoprazole or an optically active isomer (R isomer)  
thereof per one capsule;

(24) The capsule according to the above (23), wherein  
the granule according to the above (17) is filled;

(25) No. 2 to 4 capsules comprising 40 mg of an  
20 optically active isomer (R isomer) of lansoprazole per one  
capsule;

(26) The capsule according to the above (25), wherein  
the granules according to the above (17) are filled;

(27) No. 3 to 5 capsules comprising 30 mg of an  
25 optically active isomer (R isomer) of lansoprazole per

capsule;

(28) The capsule according to the above (27), wherein the granules as defined in the above (17) are filled; and

(29) A pharmaceutical composition which is a  
5 combination of an antimicrobial agent and the granules according to the above (3).

#### DETAILED DESCRIPTION OF THE INVENTION

The granules of the present invention are  
10 characterized in that they contain an acid-unstable medicament in high concentration, and have, at least, a principal ingredient layer, an intermediate coating layer which is formed on the principal ingredient layer, and further an enteric coating layer which is formed on the  
15 intermediate coating layer. The acid-unstable medicament as the principal active ingredient is contained in the principal ingredient layer but, structurally, the principal ingredient layer may have such a construction that it further has a core therein. And, the intermediate coating  
20 layer which is formed between the principal ingredient layer and the enteric coating layer has a stabilization function of the acid-unstable medicament contained in the principal ingredient layer by avoiding direct contact between the principal ingredient layer and the enteric  
25 coating layer.


---

The acid-unstable medicament in the present invention is not specifically limited, and may any medicaments which become unstable when exposed to an acid. Examples of such acid-unstable medicament include PPI having antiulcer activity, inter alia, benzimidazole compounds, imidazopyridine compounds, erythromycin antimicrobial compounds, and antiinflammatory enzymes such as serrapeptase, semialkaline proteinase, etc. and the like. In particular, the present invention is suitable for PPI such as benzimidazole compounds and imidazopyridine compounds having antiulcer activity. Hereinafter, the present invention will be illustrated with respect to benzimidazole compounds, but the present invention is not limited thereto, and can be similarly applied to other acid-unstable medicaments.

In the benzimidazole compounds having antiulcer activity used in the present invention, a preferred one is a compound of the above formula (I) in which ring A is a benzene ring which may be substituted with a halogen atom, an optionally halogenated C<sub>1-4</sub> alkyl group, an optionally halogenated C<sub>1-4</sub> alkoxy group and a 5 -or 6-membered heterocyclic group, R<sup>1</sup> is a C<sub>1-6</sub> alkyl group, a C<sub>1-6</sub> alkoxy group, a C<sub>1-6</sub> alkoxy-C<sub>1-6</sub> alkoxy group or a di-C<sub>1-6</sub> alkylamino group, R<sup>2</sup> is a hydrogen atom, a C<sub>1-6</sub> alkoxy-C<sub>1-6</sub> alkoxy group or an optionally halogenated C<sub>1-6</sub> alkoxy group, R<sup>3</sup> is a

hydrogen atom or a C<sub>1-6</sub> alkyl group, and Y is a nitrogen atom.

The particularly preferred one is a compound represented by the formula (Ia):



5 wherein R<sup>1</sup> represents a C<sub>1-3</sub> alkyl group or a C<sub>1-3</sub> alkoxy group, R<sup>2</sup> represents a C<sub>1-3</sub> alkoxy group which may be halogenated or substituted with a C<sub>1-3</sub> alkoxy group, R<sup>3</sup> represents a hydrogen atom or a C<sub>1-3</sub> alkyl group, and R<sup>4</sup> represents a hydrogen atom, an optionally halogenated C<sub>1-3</sub> alkoxy group or a pyrrolyl group (e.g. 1-, 2- or 3-pyrrolyl group).

10

In the above formula (Ia), the particularly preferred one is a compound in which R<sup>1</sup> is a C<sub>1-3</sub> alkyl group, R<sup>2</sup> is an optionally halogenated C<sub>1-3</sub> alkoxy group, R<sup>3</sup> is a hydrogen atom, and R<sup>4</sup> is a hydrogen atom or an optionally halogenated C<sub>1-3</sub> alkoxy group.

15

Examples of the "substituent" in the "optionally substituted benzene ring" represented by ring A in the compound represented by the above formula (I) [hereinafter, referred to as compound (I)] include a halogen atom, a cyano group, a nitro group, an optionally substituted alkyl

20

group, a hydroxy group, an optionally substituted alkoxy group, an aryl group, an aryloxy group, a carboxy group, an acyl group, an acyloxy group, a 5- to 10-membered heterocyclic group, etc. and the benzene ring may be substituted with about one to three these substituents. When the number of substituents is 2 or more, respective substituents may be the same or different. Among these substituents, a halogen atom, an optionally substituted alkyl group, an optionally substituted alkoxy group, etc. is preferred.

Examples of the halogen atom include fluorine, chlorine and bromine atoms, etc. Inter alia, fluorine is preferred.

Examples of the "alkyl group" in the "optionally substituted alkyl group" include C<sub>1-7</sub> alkyl groups (e.g. methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl groups, etc.). As the "substituent" in the "optionally substituted alkyl group", there can be exemplified a halogen atom, a hydroxy group, C<sub>1-6</sub> alkoxy groups (e.g. methoxy, ethoxy, propoxy, butoxy, etc.), C<sub>1-6</sub> alkoxy-carbonyl groups (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl groups, etc.), a carbamoyl group, etc. and the number of these substituents may be about 1 to 3. When the number of substituents is 2 or more, respective substituents may be

the same or different.

Examples of the "alkoxy group" in the "optionally substituted alkoxy group" include C<sub>1-6</sub> alkoxy groups (e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, pentoxy, etc.). Examples of the "substituent" in the "optionally substituted alkoxy group" include the same "substituents" as those for the above "substituent" in the "optionally substituted alkyl group", and the number of substituents is also the same as defined above.

Examples of the "aryl group" include C<sub>6-14</sub> aryl groups (e.g. phenyl, 1-naphthyl, 2-naphthyl, biphenyl, 2-anthryl groups, etc.) and the like.

Examples of the "aryloxy group" include C<sub>6-14</sub> aryloxy groups (e.g. phenyloxy, 1-naphthyloxy, 2-naphthyloxy groups, etc.) and the like.

Examples of the "acyl group" include formyl, alkylcarbonyl, alkoxycarbonyl, carbamoyl, alkylcarbamoyl, alkylsulfinyl, alkylsulfonyl groups, and the like.

Examples of the "alkylcarbonyl group" include C<sub>1-6</sub> alkyl-carbonyl groups (e.g. acetyl, propionyl groups, etc.) and the like.

Examples of the "alkoxycarbonyl group" include C<sub>1-6</sub> alkoxy-carbonyl groups (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl groups, etc.) and the like.



Examples of the "alkylcarbamoyl group" include N-C<sub>1-6</sub> alkyl-carbamoyl groups (e.g. methylcarbamoyl, ethylcarbamoyl groups, etc.), N,N-di-C<sub>1-6</sub> alkyl-carbamoyl groups (e.g. N,N-dimethylcarbamoyl, N,N-diethylcarbamoyl groups, etc.) and the like.

Examples of the "alkylsulfinyl group" include C<sub>1-7</sub> alkylsulfinyl groups (e.g. methylsulfinyl, ethylsulfinyl, propylsulfinyl, isopropylsulfinyl groups, etc.).

Examples of the "alkylsulfonyl group" include C<sub>1-7</sub> alkylsulfonyl groups (e.g. methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl groups, etc.).

Examples of the "acyloxy group" include an alkylcarbonyloxy group; an alkoxycarbonyloxy group, a carbamoyloxy group, an alkylcarbamoyloxy group, an alkylsulfinyloxy group, an alkylsulfonyloxy group and the like.

Examples of the "alkylcarbonyloxy group" include C<sub>1-6</sub> alkyl-carbonyloxy groups (e.g. acetyloxy, propionyloxy groups, etc.) and the like.

Examples of the "alkoxycarbonyloxy group" include C<sub>1-6</sub> alkoxy-carbonyloxy groups (e.g. methoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, butoxycarbonyloxy groups, etc.) and the like.

Examples of the "alkylcarbamoyloxy group" include C<sub>1-6</sub> alkyl-carbamoyloxy groups (e.g. methylcarbamoyloxy,

ethylcarbamoyloxy groups, etc.) and the like.

Examples of the "alkylsulfinyloxy group" include C<sub>1-7</sub> alkylsulfinyloxy groups (e.g. methylsulfinyloxy, ethylsulfinyloxy, propylsulfinyloxy, isopropylsulfinyloxy groups, etc.) and the like.

Examples of the "alkylsulfonyloxy group" include C<sub>1-7</sub> alkylsulfonyloxy groups (e.g. methylsulfonyloxy, ethylsulfonyloxy, propylsulfonyloxy, isopropylsulfonyloxy groups, etc.).

Examples of the "5- to 10-membered heterocyclic group" include 5- to 10-membered (preferably 5- or 6-membered) heterocyclic groups containing 1 or more (for example, 1 to 3) hetero atom(s) selected from a nitrogen atom, a sulfur atom and an oxygen atom in addition to carbon atoms, and specific examples thereof include a 2- or 3-thienyl group, a 2-, 3- or 4-pyridyl group, a 2- or 3-furyl group, a 1-, 2- or 3-pyrrolyl group, a 2-, 3-, 4-, 5- or 8-quinolyl group, a 1-, 3-, 4- or 5-isoquinolyl group, and a 1-, 2- or 3-indolyl group and the like. Among them, a preferred one is a 5- or 6-membered heterocyclic group such as a 1-, 2- or 3-pyrrolyl group, etc.

Preferably, ring A is a benzene ring optionally substituted with 1 or 2 substituent(s) selected from a halogen atom, an optionally halogenated C<sub>1-4</sub> alkyl group, an optionally halogenated C<sub>1-4</sub> alkoxy group and a 5- or 6-

membered heterocyclic group.

Examples of the "optionally substituted alkyl group" represented by  $R^1$ ,  $R^2$  or  $R^3$  include the above "optionally substituted alkyl group" described as the substituent for the above ring A.

Examples of the "optionally substituted alkoxy group" represented by  $R^1$ ,  $R^2$  or  $R^3$  include the above "optionally substituted alkoxy group" described as the substituent for the above ring A.

Examples of the "optionally substituted amino group" represented by  $R^1$ ,  $R^2$  or  $R^3$  include an amino group, mono- $C_{1-6}$  alkylamino groups (e.g. methylamino, ethylamino, etc.), mono- $C_{6-14}$  arylamino groups (e.g. phenylamino, 1-naphthylamino, 2-naphthylamino, etc.), di- $C_{1-6}$  alkylamino groups (e.g. dimethylamino, diethylamino, etc.), di- $C_{6-14}$  arylamino groups (e.g. diphenylamino, etc.) and the like.

Preferred  $R^1$  is a  $C_{1-6}$  alkyl group, a  $C_{1-6}$  alkoxy group, a  $C_{1-6}$  alkoxy- $C_{1-6}$  alkoxy group, or a di- $C_{1-6}$  alkylamino group. More preferred  $R^2$  is a  $C_{1-3}$  alkyl group or a  $C_{1-3}$  alkoxy group.

Preferred  $R^2$  is a hydrogen atom, a  $C_{1-6}$  alkoxy- $C_{1-6}$  alkoxy group or an optionally halogenated  $C_{1-6}$  alkoxy group. More preferred  $R^3$  is a  $C_{1-3}$  alkoxy group which is halogenated or optionally substituted with a  $C_{1-3}$  alkoxy group.

Preferred R<sup>3</sup> is a hydrogen atom, or a C<sub>1-6</sub> alkyl group.  
More preferred R<sup>4</sup> is a hydrogen atom or a C<sub>1-3</sub> alkyl group  
(particularly a hydrogen atom).

Preferred Y is a nitrogen atom.

5        Specific examples of the compound (I) are the  
following compounds:

2-[[[3-methyl-4-(2,2,2-trifluoroethoxy)-2-  
pyridinyl)methyl]sulfinyl]-1H-imidazole (lansoprazole), 2-  
[[[3,5-dimethyl-4-methoxy-2-pyridinyl)methyl]sulfinyl]-5-  
10 methoxy-1H-benzimidazole, 2-[[[4-(3-methoxypropoxy)-3-  
methyl-2-pyridinyl)methyl]sulfinyl]-1H-benzimidazole sodium  
salt, 5-difluoromethoxy-2-[[[3,4-dimethoxy-2-  
pyridinyl)methyl]sulfinyl]-1H-benzimidazole, etc.

Among these compounds, in particular, lansoprazole,  
15 that is, 2-[[[3-methyl-4-(2,2,2-trifluoroethoxy)-2-  
pyridinyl)methyl]sulfinyl]-1H-benzimidazole is preferred.

Examples of the imidazopyridine compound include  
tenatoprazole and the like.

The above compound (I) and imidazopyridine compound  
20 may be a racemic modification or optically active isomers  
such as R-isomer and S-isomer. For example, optically  
active isomers such as (R)-2-[[[3-methyl-4-(2,2,2-  
trifluoroethoxy)-2-pyridinyl)methyl]sulfinyl]-1H-  
benzimidazole (hereinafter, referred to as lansoprazole R  
25 isomer in some cases) and (S)-2-[[[3-methyl-4-(2,2,2-

trifluoroethoxy)-2-pyridinyl]methyl]sulfinyl]-1H-benzimidazole (hereinafter, referred to as lansoprazole S isomer in some cases) are particularly suitable in the present invention. Usually, lansoprazole, lansoprazole R isomer, lansoprazole S isomer, etc. are preferably in the form of crystals. However, not only crystalline compounds but also amorphous compounds can be used because these compounds are stabilized by formulating into preparations and, additionally, these compounds are further stabilized by blending a basic inorganic salt therein and providing with an intermediate coating layer.

As a salt of the compound (I), a pharmaceutically acceptable salt is preferred, and examples thereof include a salt with an inorganic base, a salt with an organic base, a salt with a basic amino acid and the like.

Preferred examples of the salt with an inorganic base include alkali metal salts such as a sodium salt and a potassium salt; alkaline earth metal salts such as a calcium salt and a magnesium salt; an ammonium salt and the like.

Preferred examples of the salt with an organic base include salts with alkylamines (trimethylamine, triethylamine, etc.), heterocyclic amines (pyridine, picoline, etc.), alkanolamines (ethanolamine, diethanolamine, triethanolamine, etc.), dicyclohexylamine,

N,N'-dibenzylethylenediamine and the like.

Preferred examples of the salt with a basic amino acid include salts with arginine, lysine, ornithine and the like.

Among these salts, an alkali metal salt or an alkaline  
5 earth metal salt is preferred. Inter alia, a sodium salt is preferred.

The compound (I) can be prepared by known per se methods, for example, the methods described in JP 61-50978 A, U.S. Patent No. 4,628,098, JP 10-195068 A, WO 98/21201,  
10 JP 52-62275 A, JP 54-141783 A and the like, or similar methods. The optically active compound (I) can be obtained by an optical resolving method (fractionating  
recrystallization method, chiral column method, diastereomer method, method using microorganisms or enzymes and the  
15 like), asymmetric oxidation method and the like. Lansoprazole R isomer can also be prepared by a process described, for example, in WO 00-78745, WO 01/83473 and the like.

As the benzimidazole compound having antiulcer  
20 activity used in the present invention, lansoprazole, omeprazole, rabeprazole, pantoprazole, leminoprazole, tenatoprazole (TU-199) and the like and optically active isomers thereof as well as pharmaceutically acceptable salts are preferred, and lansoprazole or an optically  
25 active isomer thereof, in particular, an R isomer is more

preferred.

The amount of PPI to be formulated in the present invention varies depending on the kind of active component and dosage, and is for example, about 12% by weight to about 40% by weight, preferably about 12% by weight to about 20% by weight, more preferably about 14% by weight to about 20% by weight based on the total granules of the present invention. When PPI is the benzimidazole compound, in particular lansoprazole, the amount is about 14% by weight to about 20% by weight.

Examples of the basic inorganic salt used in the present invention include basic inorganic salts of sodium, potassium, magnesium and calcium. A preferred one is a basic inorganic salt of magnesium or calcium. A more preferred one is a basic inorganic salt of magnesium.

Examples of the basic inorganic salt of sodium include sodium carbonate, sodium bicarbonate, sodium hydroxide and the like.

Examples of the basic inorganic salt of potassium include potassium carbonate, potassium bicarbonate, potassium hydroxide and the like.

Examples of the basic inorganic salt of magnesium include ground magnesium carbonate, magnesium carbonate, magnesium oxide, magnesium hydroxide, magnesium aluminate metasilicate, magnesium silicate, magnesium aluminate,

synthetic hydrotalcite  $[\text{Mg}_6\text{Al}_2(\text{OH})_{16}\cdot\text{CO}_3\cdot 4\text{H}_2\text{O}]$  and alumina magnesium hydroxide  $[2.5\text{MgO}\cdot\text{Al}_2\text{O}_3\cdot x\text{H}_2\text{O}]$ , preferably, ground magnesium carbonate, magnesium carbonate, magnesium oxide, magnesium hydroxide and the like.

5        Examples of the basic inorganic salt of calcium include precipitated calcium carbonate, calcium hydroxide and the like.

More preferred examples of the basic inorganic salt include ground magnesium carbonate, magnesium carbonate,  
10        magnesium oxide, magnesium hydroxide and the like.

The basic inorganic salt used in the present invention may be a salt whose 1% aqueous solution or suspension has basic pH (pH 7 or higher).

The basic inorganic salt may be formulated alone or in  
15        combination of two or more thereof, and the amount to be formulated is about 0.2 to about 0.6 part by weight, preferably about 0.2 to about 0.4 part by weight based on 1 part by weight of PPI (benzimidazole compound, etc.).

Inter alia, when PPI is lansoprazole or an optically active  
20        isomer thereof, preferably, the basic inorganic salt (preferably basic inorganic salt of magnesium or calcium, more preferably magnesium carbonate) is formulated in an amount of about 0.2 to about 0.4 part by weight based on 1 part by weight of PPI.

25        In the present invention, the granules containing PPI



(benzimidazole compound, etc.) in an amount of about 12% by weight to about 40% by weight based on the total granules, containing the basic inorganic salt as a stabilizing agent, and having an average particle size of about 600  $\mu\text{m}$  or more" are preferred. When the particle size is smaller, it is difficult to increase concentration of the benzimidazole compound because a surface area becomes larger, which requires a larger amount of an enteric layer or agent. That is, in the present invention, it has been possible to decrease the amount of an enteric layer or agent by setting the particle size of at least about 600  $\mu\text{m}$  or more, thereby increasing the concentration of the benzimidazole compound. The average particle size is about 600 to about 2500  $\mu\text{m}$ . The more preferred average particle size is about 1000 to about 2000  $\mu\text{m}$ . The granules may contain particles having a particle size of about 400 to about 3000  $\mu\text{m}$ , preferably about 500 to about 2500  $\mu\text{m}$  in so far as their average particle size, as a whole, is within the above range.

The particle size is measured by using a sieving method (Powder-Theory and Application-, p475, 1979, Maruzen), and the average particle size is calculated based on an average of meshes of corresponding sieves and weight distribution. That is, arithmetic averaging is performed based on a product of an average and each weight.

The granules of the present invention can be prepared

by a known granulation method. Examples of the granulation method include a rotary granulation method (e.g. centrifugal Fluid-bed granulation method), a fluidized granulation method, an agitation granulation method (e.g. 5 agitator fluidized granulation method) and the like. Among them, a rotary granulation method and an agitation granulation method (agitator fluidized granulation method) are preferred.

Specific examples of the rotary granulation method 10 include CF apparatus manufactured by Freund, etc. Specific examples of the agitator fluidized granulation method include methods using Spiral Flow manufactured by Freund, Multiplex manufactured by Powlex, New Malume manufactured by Fuji Powdal and the like. A method for spraying a 15 binder solution can be appropriately selected according to the kind of a granulator and, for example, it may be any of a top spraying manner, a bottom spraying manner, a tangential spraying manner and the like.

Preferably, as described above, the granules of the 20 present invention have a principal ingredient layer containing a principal ingredient, an intermediate coating layer which is formed on the principal ingredient layer, and an enteric coating layer which is formed on the intermediate coating layer.

25 For obtaining the granules having higher sphericity

and narrower particle size distribution, in the present invention, the principal ingredient layer is formed by coating cores composed of one or more materials selected from sucrose, starch, lactose and microcrystalline cellulose with the benzimidazole compound. For example, granules having a core may be prepared by the method described in JP 63-301816 A. Such granules can be obtained by a method of coating a sugar core with a powdery spreading mixture containing the benzimidazole compound having antiulcer activity, the basic metal salt, an excipient, a disintegrating agent and the like, while spraying a binder solution of hydroxypropylcellulose on the sugar cores. Examples of the core granules include Nonpareil obtained by coating sucrose (75 parts by weight) with corn starch (25 parts by weight) by a known per se method, spherical core granules using microcrystalline cellulose, etc. Alternatively, core granules per se may be an active ingredient which becomes the above principal ingredient layer. An average particle size of the core granules is generally 14 to 80 mesh.

Examples of the cores include a spherical granulated material of sucrose and starch, a spherical granulated material of crystalline cellulose, a spherical granulated material of crystalline cellulose and lactose, etc.

It is desirable that cores are as uniformly spherical

as possible so as to reduce variability of coating.

The proportion of the coating layer to the cores can be selected from within such a range that the dissolution property of the benzimidazole compound and a particle size of the granules can be controlled. For example, the proportion is usually about 0.2 part by weight to about 5 parts by weight, preferably about 0.1 part by weight to about 5 parts by weight based on 1 part by weight of the cores.

Coating layers with which the principal ingredient layer is coated may be formed of plural layers. The plural coating layers may contain various coating layers such as a coating layer for subcoating in addition to the intermediate coating layer containing no medicament and the enteric coating layer, and a particular combination of those coating layers may be appropriately selected.

In enteric coating granules containing an unstable principal ingredient such as the benzimidazole compound, etc., from a viewpoint of improvement in stability of the principal ingredient, it is preferable to arrange an intermediate coating layer between a principal ingredient layer containing the benzimidazole compound, etc. and an enteric coating layer to block direct contact between the layers because the enteric coating layer component is an acidic substance.

Such an intermediate coating layer may be a coating layer which can prevent contact between the benzimidazole compound as a basis and an enteric coating layer, and the amount and material of the coating layer are not limited in so far as such an objective is achieved. For example, there is a layer in which a saccharide such as sucrose [refined white sugar (pulverized (powdered sugar) or not pulverized), etc.], starch sugar such as corn starch, lactose, honey, sugar alcohol (D-mannitol, erythritol, etc.), etc. is appropriately formulated into a polymer base such as low-substituted hydroxypropylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose (e.g. TC-5, etc.), polyvinylpyrrolidone, polyvinyl alcohol, methylcellulose, hydroxyethylmethylcellulose and the like. In addition, to the intermediate coating layer may be appropriately added an excipient (e.g. masking agent (titanium oxide, etc.), and an antistatic agent (titanium oxide, talc, etc.)) which are added for formulating into a preparation as needed, as described hereinafter.

The amount of the intermediate coating layer to be coated is usually about 0.02 part by weight to about 1.5 parts by weight, preferably about 0.05 part by weight to about 1 part by weight based on 1 part by weight of the granules containing, for example, benzimidazole. Coating can be performed according to a conventional method. For

example, preferably, these intermediate coating layer components are diluted with purified water or the like to obtain a liquid, which is sprayed for coating. At this time, it is preferable to perform coating while spraying a binder agent of hydroxypropylcellulose or the like.

Examples of the "enteric coating layer" to be used for coating the granules in the present invention include aqueous enteric polymer bases such as cellulose acetate phthalate (CAP), hydroxypropylmethylcellulose phthalate, hydroxymethylcellulose acetate succinate, methacrylic acid copolymer, carboxymethylethylcellulose, shellac and the like, sustained-release bases such as ethyl acrylate-methacrylic acid copolymer and the like, and plasticizers such as water-soluble polymer, triethyl citrate, polyethylene glycol, acetylated monoglyceride, triacetin, castor oil and the like. They can be used alone or in combination of two or more thereof.

The enteric coating layer is an enteric polymer base, preferably an aqueous enteric methacrylic acid copolymer.

The amount of the enteric coating layer to be coated is about 10% by weight to about 70% by weight, preferably about 10% by weight to about 50% by weight, more preferably about 15% by weight to about 30% by weight based on the total amount of the granules before coating of the enteric coating.

Further, additives for formulating into pharmaceutical preparations can be used, and examples thereof include excipients (e.g. glucose, fructose, lactose, sucrose, D-mannitol, erythritol, maltitol, trehalose, sorbitol, corn starch, potato starch, wheat starch, rice starch, crystalline cellulose, anhydrous silicic acid, anhydrous calcium phosphate, precipitated calcium carbonate, calcium silicate, etc.), binders (e.g. hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, methylcellulose, polyvinyl alcohol, sodium carboxymethylcellulose, partial  $\alpha$ -starch,  $\alpha$ -starch, sodium alginate, pullulan, gum arabic powder, gelatin etc.), disintegrating agents (e.g. low-substituted hydroxypropylcellulose, carmellose, potassium carmellose, sodium carboxymethylstarch, sodium crosscarmellose, crosspovidone, hydroxypropylstarch, etc.), corrigents (e.g. citric acid, ascorbic acid, tartaric acid, malic acid, aspartame, potassium acesulfame, sormatin, saccharin sodium, dipotassium glycyrrhizin, sodium glutamate, sodium 5'-inosinate, sodium 5'-guanylate, etc.), surfactants (e.g. polysorbates (polysorbate 80, etc.), polyoxyethylene-polyoxypropylene copolymer, sodium laurylsulfate, etc.), flavors (e.g. lemon oil, orange oil, menthol, mint oil, etc.), lubricants (e.g. magnesium stearate, sucrose fatty acid esters, sodium stearyl fumarate, stearic acid, talc,

polyethylene glycol, etc.), colorants (e.g. titanium oxide, edible Yellow No.5, edible Blue No.2, iron sesquioxide, yellow iron sesquioxide etc.), antioxidants (e.g. sodium ascorbate, L-cysteine, sodium sulfite, etc.), masking  
5 agents (e.g. titanium oxide, etc.), antistatic agents (e.g. talc, titanium oxide, etc.) and the like.

The particle size of raw materials used for these materials is not particularly limited, but particles of about 500  $\mu\text{m}$  or smaller are preferred from a viewpoint of  
10 manufacturing properties and administration properties.

The granules of the present invention can also be used in granular preparations, capsules, tablets, effervescent preparations, suspensions or the like.

From a viewpoint of easy handling, etc., capsules and  
15 tablets are preferred. As capsules, gelatin capsules, HPMC capsules, pullulan capsules and the like may be used. When used as capsules, capsules of No.3 to No.5 in size are preferable for easy administration. For example, in case of a capsule containing lansoprazole-containing granules,  
20 preferably, granules having an average particle size of about 1000  $\mu\text{m}$  to about 2000  $\mu\text{m}$  are prepared by providing an intermediate coating layer on a principal ingredient layer, which contains lansoprazole in an amount of about 14% by weight to about 20% by weight based on the total granules  
25 and a basic salt of magnesium and/or calcium in an amount



of about 0.2 part by weight to about 0.4 part by weight based on 1 part by weight of lansoprazole, and then providing an enteric coating layer thereon, and the granules are filled in a capsule. As a capsule containing 5 30 mg of lansoprazole per capsule, conventional products are No.1 to No.2 capsules, while No.3 to No.5 stable capsules can be produced according to the present invention. In addition, in case of a capsule containing 15 mg of lansoprazole per capsule in which the above granules are 10 filled, it is possible to reduce the size to No.4 to No.5 capsules. Further, in case of a capsule containing 60 mg of lansoprazole R isomer, No.3 to No. 1 capsules are possible. Furthermore, in case of a capsule containing 40 mg, No.4 to No.2 capsules are possible and, in case of a 15 capsule containing 30 mg, No.5 to No.3 capsules are possible.

In the granules of the present invention, since PPI such as the benzimidazole compound has excellent antiulcer activity, gastric acid secretion inhibiting activity, 20 mucosa protecting activity, anti-*Helicobacter pylori* activity and the like, and has low toxicity, they are useful for medicine. In this case, the granules of the present invention can be administered to a mammal (e.g. human, monkey, sheep, horse, dog, cat, rabbit, rat, mouse, 25 etc.) orally for the purpose of treating and preventing

peptic ulcers (e.g. gastric ulcer, duodenal ulcer, stomal  
ulcer, etc.), Zollinger-Ellison syndromes, gastritis,  
reflux esophagitis, symptomatic gastroesophageal reflux  
disease (symptomatic GERD)), NUD (non ulcer dyspepsia),  
5 stomach cancer (including stomach cancer accompanied with  
promotion of production of interleukin-1 $\beta$  due to genetic  
polymorphism of interleukin-1), gastric MALT lymphoma and  
the like, eradicating *Helicobacter pylori*, suppressing  
upper gastrointestinal hemorrhage due to peptic ulcers,  
10 acute stress ulcer and hemorrhagic gastritis, suppressing  
upper digestive tract hemorrhagic due to invasion stress  
(stress resulting from major operation requiring post-  
operative intensive management, and cerebrovascular  
disorder, head trauma, multiple organ failure and diffuse  
15 burn requiring intensive care), treating and preventing  
ulcers resulting from nonsteroidal antiinflammatory  
medicaments; treating and preventing excess stomach acid  
and ulcer due to post-operation stress, and the like. For  
eradicating *Helicobacter pylori*, etc., the granules or  
20 capsule of the present invention may be used together with  
other active component(s) (e.g. 1 to 3 active component(s)).

Examples of the "other active component(s)" include  
anti-*Helicobacter pylori* active substances, antimicrobial  
agents such as imidazole compounds, quinolone compounds,  
25 and bismuth salts. Inter alia, medicines comprising a

combination of the granules or capsule of the present invention and an antimicrobial agent are preferred. Among them, a combination with anti-*Helicobacter pylori* active substances, or antimicrobial agents such as imidazole compounds is preferred. Examples of the "anti-*Helicobacter pylori* active substance" include penicillin antibiotics (e.g. amoxicillin, benzylpenicillin, piperacillin, mecillinam, etc.), cephem antibiotics (e.g. cefixime, cefaclor, etc.), macrolide antibiotics (e.g. erythromycin antibiotics such as erythromycin, clarithromycin, etc.), tetracycline antibiotics (e.g. tetracycline, minocycline, streptomycin, etc.), aminoglycoside antibiotics (e.g. gentamycin, amikacin, etc.), imipenem and the like. Inter alia, penicillin antibiotics and macrolide antibiotics are preferred.

Examples of the "imidazole compound" include metronidazole, miconazole, etc. Examples of the "bismuth salt" include bismuth acetate, bismuth citrate, etc. Antibacterial agents of "quinolone compounds" are also preferred, and examples thereof include ofloxacin, ciprofloxacin, etc. Inter alia, for eradicating *Helicobacter pylori*, it is preferred to use the granules or capsule of the present invention in combination with penicillin antibiotics (e.g. amoxicillin, etc.) and/or erythromycin antibiotics (e.g. clarithromycin etc.).

For example, in case of lansoprazole, in many cases, a conventional 15 mg-containing capsule is a product filled in a No.3 capsule, and a conventional 30 mg-containing capsule is a product filled in a No.1 capsule. However, according to the present invention, since the amounts of components other than a basis can be reduced without deteriorating stability of the basis and a preparation, the size of a 15 mg-containing capsule can be reduced to No.4 to No.5 capsules, and the size of a 30 mg-containing capsule can be reduced to No.3 to No.5 capsules, respectively.

Further, even in a 60 mg-containing capsule, it is possible to use No.1 to No.3 capsules.

Furthermore, in case of an optically active isomer of lansoprazole, No.3 to No.5 capsules, No.2 to No.4 capsules and No.1 to No.3 capsules can be used for 30 mg-, 40 mg- and 60 mg-containing capsules, respectively.

For example, because of a high content of an active component and easy administration, a capsule containing 60 mg of lansoprazole or lansoprazole R isomer is suitable for treating, inter alia, acid hypersecretion symptom including Zollinger-Ellison syndromes.

A daily dosage varies depending on the degree of symptom, the age, sex and weight of subject, the administration time, interval, and the kind of principal or

active ingredient, etc., and is not specifically limited. For example, when orally administered to an adult (60 kg) as an antiulcer agent, a dosage is about 0.5 to 1500 mg/day, preferably about 5 to 150 mg/day in terms of an active  
5 component. These benzimidazole compound-containing preparations may be administered once or by dividing into 2 to 3 times daily.

Hereinafter, the present invention will be illustrated in more detail by Examples, but the present invention is  
10 not limited by them. In the following Examples, lansoprazole and its optically active isomer in the form of crystals were used.

#### Example 1

15       Lansoprazole, magnesium carbonate, sucrose (pulverized sucrose), corn starch and low-substituted hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture of principal ingredient. Spherical granules consisting of sucrose and starch were placed in an  
20 agitator fluidized granulation coating machine (MP-10 manufactured by Powlex), and the above spreading mixture of principal ingredient was coated while spraying a hydroxypropylcellulose solution (2.5%: W/W) to obtain spherical granules. The resultant spherical granules were  
25 dried in a vacuum at 40°C for 16 hours, and passed through

a round sieve to obtain granules of 500  $\mu\text{m}$  to 1180  $\mu\text{m}$ .

Granules Composition in 150 mg

Spherical granules consisting of sucrose and starch	50 mg
Lansoprazole	30 mg
Magnesium carbonate	10 mg
Sucrose (pulverized sucrose)	30 mg
Corn starch	14 mg
Low-substituted hydroxypropylcellulose	15 mg
Hydroxypropylcellulose	1 mg
Purified water	39 $\mu\text{l}$
Total	150 mg

5 An enteric suspension having the following composition was coated on the above granules using an agitator fluidized granulation coating machine (MP-10, manufactured by Powlex), which were dried as such, and passed through a round sieve to obtain enteric granules of 710 to 1420  $\mu\text{m}$ .

10 Into the resultant granules were mixed talc and aerosil, and 190 mg of the resultant mixed granules were filled into a No.3 capsule manually.

Composition of enteric suspension

Methacrylic acid copolymer	86.7 mg (solid components 26 mg)
Talc	7.8 mg
Polyethylene glycol	2.5 mg
Titanium oxide	2.5 mg
Polysorbate 80	1.0 mg
Purified water	119.5 $\mu\text{l}$
Total	39.8 mg (as solids)

## Composition of enteric granules

Granules	150 mg
Enteric coating layer	39.8 mg
Total	189.8 mg

## Composition of mixed granules

Enteric granules	189.8 mg
Talc	0.1 mg
Aerosil	0.1 mg
Total	190 mg

## 5 Composition of capsule

Mixed granules	190 mg
No. 3 capsule	1

In addition, particle size distribution of the resultant mixed granules was measured using a round sieve, and the results are shown below:

1180 $\mu$ m remaining	10.6%
1180/1000 $\mu$ m	70.9%
1000/850 $\mu$ m	12.0%
850 $\mu$ m pass	6.4%

10

## Example 2

Lansoprazole, magnesium carbonate, sucrose (pulverized sucrose), corn starch and low-substituted hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture of principle ingredient. Sucrose (pulverized sucrose), corn starch and low-substituted hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture for intermediate layer. Spherical

15

granules consisting of sucrose and starch were placed in a centrifugal Fluid-bed granulator (CF-360 $\phi$  manufactured by Freund), and the above spreading mixture of principal ingredient and the spreading mixture for intermediate layer were coated while spraying a hydroxypropylcellulose solution (2.5%: W/W) to obtain spherical granules. The resultant spherical granules were dried in a vacuum at 40°C for 16 hours, and passed through a round sieve to obtain granules of 500  $\mu$ m to 1180  $\mu$ m.

10

## Granules Composition in 160 mg

Spherical granules consisting of sucrose and starch	50 mg
Spreading mixture of principal ingredient	
Lansoprazole	30 mg
Magnesium carbonate	10 mg
Sucrose (pulverized sucrose)	30 mg
Corn starch	14 mg
Low-substituted hydroxypropylcellulose	15 mg
Spreading mixture for intermediate layer	
Sucrose (pulverized sucrose)	5 mg
Corn starch	2.5 mg
Low-substituted hydroxypropylcellulose	2.5 mg
Binder solution	
Hydroxypropylcellulose	1 mg
Purified water	34 $\mu$ l
Total	160 mg

An enteric suspension having the following composition was coated on the above granules using an agitator fluidized granulation coating machine (MP-10 manufactured

15



by Powlex), which was dried as such, and passed through a round sieve to obtain enteric granules of 600 to 1420  $\mu\text{m}$ . Into the resultant granules were mixed talc and aerosil, and 200 mg of the resultant mixed granules were filled into a No.3 capsule.

## Composition of enteric suspension

Methacrylic acid copolymer	86.7 mg (solid components
	26 mg)
Talc	7.8 mg
Polyethylene glycol	2.5 mg
Titanium oxide	2.5 mg
Polysorbate 80	1.0 mg
Purified water	119.5 $\mu\text{l}$
Total	39.8 mg (as solids)

## Composition of enteric granules

Granules	160 mg
Enteric coating layer	39.8 mg
Total	199.8 mg

## Composition of mixed granules

Enteric granules	199.8 mg
Talc	0.1 mg
Aerosil	0.1 mg
Total	200 mg

## Composition of capsule

Mixed granules	200 mg
No.3 capsule	1

In addition, particle size distribution of the resultant mixed granules was measured using a round sieve,

and the results are shown below:

1180 $\mu$ m remaining	20.2%
1180/1000 $\mu$ m	76.2%
1000/850 $\mu$ m	3.6%
850/710 $\mu$ m	0.0%
710 $\mu$ m passed	0.0%

### Example 3

A suspension was prepared by using lansoprazole,  
5 magnesium carbonate, low-substituted hydroxypropylcellulose,  
hydroxypropylcellulose and purified water.  
Microcrystalline cellulose spherical granules were placed  
in an agitator fluidized granulation coating machine (MP-10  
manufactured by Powlex), and the suspension was coated  
10 while spraying the suspension to obtain spherical granules.  
The granules were dried as such, and passed through a round  
sieve to obtain granules of 500  $\mu$ m to 1180  $\mu$ m.

### Granules Composition in 70 mg

Microcrystalline cellulose spherical granules	20 mg
Lansoprazole	30 mg
Magnesium carbonate	10 mg
Low-substituted hydroxypropylcellulose	5 mg
Hydroxypropylcellulose	5 mg
Purified water	100 $\mu$ l
Total	70 mg

15

An intermediate layer suspension was prepared by using  
hydroxypropylmethylcellulose, low-substituted  
hydroxypropylcellulose, D-mannitol and purified water. The

granules containing lansoprazole were placed in an agitator fluidized granulation coating machine (MP-10 manufactured by Powlex), and an intermediate layer was coated while spraying the suspension to obtain spherical granules. The

5. granules were dried as such and passed through a round sieve to obtain granules of 500  $\mu\text{m}$  to 1800  $\mu\text{m}$ .

Granules Composition in 80 mg

Lansoprazole-containing granules	70 mg
Hydroxypropylmethylcellulose	5 mg
Low-substituted hydroxypropylcellulose	2.5 mg
D-mannitol	2.5 mg
Purified water	40 $\mu\text{l}$
Total	80 mg

An enteric suspension having the following composition

10 was coated on the above granules using an agitator fluidized granulation coating machine (MP-10 manufactured by Powlex); which was dried as such, and passed through a round sieve to obtain enteric granules of 600 to 1420  $\mu\text{m}$ . Into the resultant granules were mixed talc and aerosil,

15 and 100 mg of the resultant mixed granules were filled in a No. 5 capsule.

Composition of enteric suspension

Methacrylic acid copolymer	43.3 mg (solid components 13 mg)
Talc	3.8 mg
Polyethylene glycol	1.2 mg
Titanium oxide	1.2 mg
Polysorbate 80	0.5 mg
Purified water	60 $\mu\text{l}$
Total	19.7 mg (as solids)

## Composition of enteric granules

Granules	80 mg
Enteric coating layer	19.7 mg
Total	99.7 mg

## Composition of mixed granules

Enteric granules	99.7 mg
Talc	0.2 mg
Aerosil	0.1 mg
Total	100 mg

## 5 Composition of capsule

Mixed granules	100 mg
No. 5 capsule	1

In addition, particle size distribution of the resultant mixed granules was measured using a round sieve, and the results are shown below:

1180 $\mu$ m remaining	5.6%
1180/1000 $\mu$ m	91.3%
1000/850 $\mu$ m	3.1%
850 $\mu$ m passed	0.0%

10

## Example 4

Lansoprazole, magnesium carbonate, sucrose (pulverized sucrose), corn starch, low-substituted hydroxypropylcellulose and hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture of principal ingredient. Sucrose (pulverized sucrose), corn starch, low-substituted hydroxypropylcellulose and hydroxypropylcellulose were thoroughly mixed to obtain a

spreading mixture for intermediate layer. Spherical granules consisting of sucrose and starch were placed in a centrifugal Fluid-bed granulator (CF-1300 $\phi$  manufactured by Freund) and the above spreading mixture of principal  
 5 ingredient and the spreading mixture for intermediate layer were coated while spraying purified water to obtain spherical granules. The resultant spherical granules were dried in a vacuum at 45°C for 18 hours, and passed through a vibrating screen to obtain granules of 500  $\mu$ m to 1180  $\mu$ m.

10

## Granules Composition in 320 mg

Spherical granules consisting of sucrose and starch	100 mg
Spreading mixture of principal ingredient	
Lansoprazole	60 mg
Magnesium carbonate	20 mg
Sucrose (pulverized sucrose)	60 mg
Corn starch	28 mg
Low-substituted hydroxypropylcellulose	30 mg
Hydroxypropylcellulose	1.8 mg
Spreading mixture for intermediate layer	
Sucrose (pulverized sucrose)	10 mg
Hydroxypropylcellulose	0.2 mg
Purified water	60 $\mu$ l
Total	320 mg

An enteric suspension having the following composition was coated on the above granules using a fluidized bed  
 15 coating machine (FLO-90 manufactured by Freund), which was dried as such, and passed through a vibrating screen to obtain enteric granules of 600 to 1420  $\mu$ m. To the

resultant granule were mixed talc and aerosil using a  
tumbler mixer (manufactured by Showa Kagakukikai-kousakusho,  
1300L), and 400 mg of the resultant mixed granules were  
filled in a No. 1 capsule using a capsule filler (MATIC-90  
5 manufactured by IMA).

## Composition of enteric suspension

Methacrylic acid copolymer	173.4 mg (solid components 52 mg)
Talc	15.6 mg
Polyethylene glycol	5.0 mg
Titanium oxide	5.0 mg
Polysorbate 80	2.0 mg
Purified water	239 $\mu$ l
Total	79.6 mg (as solids)

## Composition of enteric granules

Granules	320 mg
Enteric coating layer	79.6 mg
Total	399.6 mg

10

## Composition of mixed granules

Enteric granules	399.6 mg
Talc	0.2 mg
Aerosil	0.2 mg
Total	400 mg

## Composition of capsule

Mixed granules	400 mg
No. 1 capsule	1

15

In addition, particle size distribution of the  
resultant mixed granules was measured using a round sieve,

and the results are shown below:

1180 $\mu\text{m}$ remaining	2.6%
1180/1000 $\mu\text{m}$	92.2%
1000/850 $\mu\text{m}$	4.6%
850/710 $\mu\text{m}$	0.4%
710 $\mu\text{m}$ passed	0.2%

#### Example 5

A composition is shown in Table 1. Lansoprazole R isomer, magnesium carbonate, sucrose (pulverized sucrose), corn starch and low-substituted hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture of principal ingredient. In addition, sucrose (pulverized sucrose), corn starch and low-substituted hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture of intermediate layer. Spherical granulates consisting of sucrose and starch were placed in a centrifugal Fluid-bed granulator (CF manufactured by Freund), and the above spreading mixture of principal ingredient and the spreading mixture for intermediate layer were successively coated while spraying a hydroxypropylcellulose solution (2%: W/W) to obtain spherical granules. The coating operation conditions were as follows: rotor rotating speed: 300 rpm, spray rate: 1.8 g/min, spray air pressure: 0.2 kg/cm<sup>2</sup>, and slit air pressure: 0.2 kg/cm<sup>2</sup>. The resultant spherical granules were dried in a vacuum at 40°C for 20 hours and passed through a round sieve to obtain granules of 710  $\mu\text{m}$

to 1420  $\mu\text{m}$ .

An enteric suspension was coated on the above granules using a fluidized granulation coating machine (LAB-1 manufactured by Powlex), which was dried as such, and passed through a round sieve to obtain enteric granules of 850 to 1420  $\mu\text{m}$ . The coating operation conditions were as follows: inlet air volume: 0.6  $\text{m}^3/\text{min}$ , inlet air temperature: 85°C, spray rate: 8 g/min, and spray air pressure: 1  $\text{kg}/\text{cm}^2$ .

Into the resultant granules were mixed talc and aerosil, and 150 mg (corresponding to 30 mg of lansoprazole R isomer), 200 mg (corresponding to 40 mg of lansoprazole R isomer) and 300 mg (corresponding to 60 mg of lansoprazole R isomer) of the resultant mixed granules were filled in No. 4, No. 3 and No. 2 capsules, respectively.

#### Example 6

A composition is shown in Table 1. Lansoprazole R isomer, magnesium carbonate, sucrose (pulverized sucrose) and low-substituted hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture of principal ingredient. In addition, sucrose (pulverized sucrose), low-substituted hydroxypropylcellulose and titanium oxide were thoroughly mixed to obtain a spreading mixture for intermediate layer. Spherical granules consisting of sucrose and starch were



placed in a centrifugal Fluid-bed granulator (CF  
manufactured by Freund), the above spreading mixture of  
principal ingredient and the spreading mixture for  
intermediate layer were successively coated while spraying  
5 a hydroxypropylcellulose solution (2%: W/W) to obtain  
spherical granules. The coating operation conditions were  
as follows: rotor rotating speed: 300 rpm, spray rate: 1.8  
g/min, spray air pressure: 0.2 kg/cm<sup>2</sup>, and slit air  
pressure: 0.2 kg/cm<sup>2</sup>. The resultant spherical granules  
10 were dried in a vacuum at 40°C for 20 hours, and passed  
through a round sieve to obtain granules of 710 µm to 1420  
µm.

An enteric suspension was coated on the above granules  
using a fluidized granulation coating machine (LAB-1  
15 manufactured by Powlex), which was dried as such, and  
passed through a round sieve to obtain enteric granules of  
850 to 1420 µm. The coating operation conditions were as  
follows: inlet air volume: 0.6 m<sup>3</sup>/min, inlet air  
temperature: 85°C, spray rate: 8 g/min, and spray air  
20 pressure: 1 kg/cm<sup>2</sup>.

Into the resultant granules were mixed talc and  
aerosil, and 150 mg (corresponding to 30 mg of lansoprazole  
R isomer), 200 mg (corresponding to 40 mg of lansoprazole R  
isomer) and 300 mg (corresponding to 60 mg of lansoprazole  
25 R isomer) of the resultant mixed granules were filled in No.

4, No. 3 and No. 2 capsules, respectively.

#### Example 7

A composition is shown in Table 1. Lansoprazole R isomer, magnesium carbonate, sucrose (pulverized sucrose), low-substituted hydroxypropylcellulose and titanium oxide were thoroughly mixed to obtain a spreading mixture of principal ingredient. Spherical granules consisting of sucrose and starch were placed in a centrifugal Fluid-bed granulator (CF manufactured by Freund), and the above spreading mixture of principal ingredient was coated while spraying a hydroxypropylcellulose solution (2%: W/W) to obtain spherical granules. The coating operation conditions were as follows: rotor rotating speed: 300 rpm, spray rate: 1.8 g/min, spray air pressure: 0.2 kg/cm<sup>2</sup>, and slit air pressure: 0.2 kg/cm<sup>2</sup>. The resultant spherical granules were dried in a vacuum at 40°C for 20 hours, and passed through a round sieve to obtain granules of 710 µm to 1420 µm.

An enteric suspension was coated on the above granules using a fluidized granulation coating machine (LAB-1 manufactured by Powlex), which was dried as such, and passed through a round sieve to obtain enteric granules of 850 to 1420 µm. The coating operation conditions were as follows: inlet air volume: 0.6 m<sup>3</sup>/min, inlet air

temperature: 85°C, spray rate: 8 g/min, and spray air pressure: 1 kg/cm<sup>2</sup>.

Into the resultant granules were mixed talc and aerosil, and 150 mg (corresponding to 30 mg of lansoprazole R isomer), 200 mg (corresponding to 40 mg of lansoprazole R isomer) and 300 mg (corresponding to 60 mg of lansoprazole R isomer) of the resultant mixed granules were filled in No. 4, No. 3 and No. 2 capsules, respectively.

Table 1: Composition table

Granules	Composition in 160 mg	Example 5	Example 6	Example 7
Spherical granules consisting of sucrose and starch		50 mg	50 mg	50 mg
Spreading mixture of principal ingredient				
Lansoprazole R isomer	40 mg	40 mg	40 mg	
Magnesium carbonate	14 mg	14 mg	14 mg	
Sucrose (pulverized sucrose)	26 mg	26 mg	36 mg	
Corn starch	9 mg	0 mg	0 mg	
Low-substituted hydroxypropylcellulose	10 mg	10 mg	12.5 mg	
Titanium oxide	0 mg	0 mg	6.5 mg	
Spreading mixture for intermediate layer				
Sucrose (pulverized sucrose)	5 mg	10 mg		
Corn starch	2.5 mg	0 mg		
Low-substituted hydroxypropylcellulose	2.5 mg	2.5 mg		
Titanium oxide	0 mg	6.5 mg		
Binder solution				
Hydroxypropylcellulose	1 mg	1 mg	1 mg	
Purified water	49 µl	49 µl	49 µl	
Total	160 mg			

## Composition of enteric suspension

Methacrylic acid copolymer	86.7 mg (solid components 26 mg)
Talc	7.8 mg
Polyethylene glycol	2.5 mg
Titanium oxide	2.5 mg
Polysorbate 80	1.0 mg
Purified water	119.5 µl
Total	39.8 mg (as solids)

## Composition of enteric granules

Granules	160 mg
Enteric coating layer	39.8 mg
Total	199.8 mg

5

## Composition of mixed granules

Enteric granules	199.8 mg
Talc	0.1 mg
Aerosil	0.1 mg
Total	200 mg

## Composition of capsules

Lansoprazole	corresponding	corresponding	corresponding
R isomer	to 30 mg	to 40 mg	to 60 mg
Mixed granules	150 mg	200 mg	300 mg
Capsule	1 (No. 4)	1 (No. 3)	1 (No. 2)

## 10 Example 8

A composition is shown in Table 2. Lansoprazole R isomer, magnesium carbonate, sucrose (pulverized sucrose), corn starch and low-substituted hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture of principal

ingredient. In addition, sucrose (pulverized sucrose), corn starch and low-substituted hydroxypropylcellulose were thoroughly mixed to obtain a spreading mixture for overcoating. Spherical granules consisting of sucrose and starch were placed in a centrifugal Fluid-bed granulator (CF manufactured by Freund), and the above spreading mixture of principal ingredient was successively coated while spraying a hydroxypropylcellulose solution (2%: W/W) to obtain spherical granules. The coating operation conditions were as follows: rotor rotating speed: 300 rpm, spray rate: 1.8 g/min, spray air pressure: 0.2 kg/cm<sup>2</sup>, and slit air pressure: 0.2 kg/cm<sup>2</sup>. The resultant spherical granules were dried in a vacuum at 40°C for 20 hours, and passed through a round sieve to obtain granules of 710 µm to 1420 µm.

A suspension composed of hydroxypropylmethylcellulose, titanium oxide and purified water shown in Table 2 was prepared. This suspension was coated on the above granules using a fluidized granulation coating machine (LAB-1 manufactured by Powlex), which was dried as such, and passed through a round sieve to obtain enteric granules of 850 to 1420 µm. The coating operation conditions were as follows: inlet air rate: 0.8 to 1 m<sup>3</sup>/min, inlet air temperature: 85°C, spray rate: 6 g/min, and spray air pressure: 0.8 to 1 kg/cm<sup>2</sup>.

An enteric suspension was coated on the above granules using a fluidized granulation coating machine (LAB-1 manufactured by Powlex), which was dried as such, and passed through a round sieve to obtain enteric granules of 850 to 1420  $\mu\text{m}$ . The coating operation conditions were as follows: inlet air rate: 0.6  $\text{m}^3/\text{min}$ , inlet air temperature: 85°C, spray rate: 8 g/min, and spray air pressure: 1  $\text{kg}/\text{cm}^2$ .

Into the resultant granules were mixed talc and aerosil, and 150 mg (corresponding to 30 mg of lansoprazole R isomer), 200 mg (corresponding to 40 mg of lansoprazole R isomer) and 300 mg (corresponding to 60 mg of lansoprazole R isomer) of the resultant mixed granules were filled into No. 4, No. 3 and No. 2 capsules, respectively.

Table 2: Composition table

Granules Composition in 141 mg

Spherical granules consisting of sucrose and starch	50 mg
Spreading mixture of principal ingredient	
Lansoprazole R isomer	40 mg
Magnesium carbonate	14 mg
Sucrose (pulverized sucrose)	26 mg
Low-substituted hydroxypropylcellulose	10 mg
Binder solution	
Hydroxypropylcellulose	1 mg
Purified water	49 $\mu\text{l}$
Total	141 mg

## Composition of intermediate layer suspension

Hydroxypropylmethylcellulose	12.5 mg
Titanium oxide	6.5 mg
Purified water	171 µl
Total	19 mg (as solids)

## Composition of intermediate layer coated granules

Granules	141 mg
Intermediate layer coating	19 mg
Total	160 mg

## 5 Composition of enteric suspension

Methacrylic acid copolymer	86.7 mg (solid components 26 mg)
Talc	7.8 mg
Polyethylene glycol	2.5 mg
Titanium oxide	2.5 mg
Polysorbate 80	1.0 mg
Purified water	119.5 µl
Total	39.8 mg (as solids)

## Composition of enteric granules

Granules	160 mg
Enteric coating layer	39.8 mg
Total	199.8 mg

## Composition of mixed granules

Enteric granules	199.8 mg
Talc	0.1 mg
Aerosil	0.1 mg
Total	200 mg

## Composition of capsules

Lansoprazole	30 mg	40 mg	60 mg
R isomer	equivalent	equivalent	equivalent
Mixed granules	150 mg	200 mg	300 mg
Capsule	1 (No. 4)	1 (No. 3)	1 (No. 2)

## Experiment 1

Each of the enteric granules (mixed granules) prepared in Examples 1 and 2 was stored in a tightly sealed bottle under 40°C/75%RH for 24 weeks and, thereafter, a content (remaining rate) and appearance change ( $\Delta E$ ) were measured. The content was measured by HPLC method. Appearance change ( $\Delta E$ ) was obtained by measuring a color difference ( $\Delta E$ ) using SM color computer SM-5 (manufactured by Suga Shikenki). The results are shown in Table 3.

Table 3

Storing conditions	Example 1		Example 2	
	$\Delta E$	Content (remaining rate)	$\Delta E$	Content (remaining rate)
Initial	—	100%	—	100%
40°C 75%RH tightly sealed 4W	1.73	100.0%	2.84	99.7%
40°C 75%RH tightly sealed 12W	7.27	100.0%	3.79	99.9%
40°C 75%RH tightly sealed 24W	3.88	98.7%	5.06	100.4%

As a result, it was revealed that the granules of the present invention had less appearance change and stable content.



## Experiment 2

Each of the enteric granules (mixed granules) prepared in Examples 5 to 8 was stored in a tightly sealed glass bottle at 60°C for 2 weeks and, thereafter, a content (remaining rate) was measured. The content was measured by HPLC method. The results are shown in Table 4. Appearance change of the preparations of Examples 7 and 8 was hardly observed with naked eyes.

10

Table 4: Results of stability test of Examples 5 to 8  
Storing conditions: 60°C tightly sealed glass bottle, 2 weeks storage

Example	Content (remaining rate)
Example 5	97.8%
Example 6	97.1%
Example 7	95.9%
Example 8	99.1%

As a result, it was revealed that the granules of the present invention are stable from a viewpoint of the content.

## INDUSTRIAL APPLICABILITY

By optimizing a blending ratio of a basic inorganic salt with an acid-unstable medicament, in particular, a benzimidazole compound, and an average particle size,

unexpectedly, the granules of the present invention enabled the acid-unstable medicament to be stable even at high concentration and high content. In addition, since the granules of the present invention contains an active  
5 ingredient at high concentration, the amount of a whole preparation can be reduced even at the same content and, therefore, the size of a capsule or the like can be reduced, and a preparation which can be easily administered can be obtained. As a result, the preparation can be easily  
10 administered to patients, in particular, elderly and pediatric patients who have difficulty in swallowing, and compliance can be improved.


## CLAIMS

1. Granules comprising a principal ingredient layer containing an acid-unstable medicament in an amount of about 12% by weight or more based on the total granules; an intermediate coating layer which is formed on the principal ingredient layer; and an enteric coating layer which is formed on the intermediate coating layer; wherein said granules contain a basic inorganic salt, and have an average particle size of about 600  $\mu\text{m}$  or more.

2. The granules according to claim 1, wherein the basic inorganic salt is a magnesium salt or a calcium salt.

3. The granules according to claim 1, wherein the acid-unstable medicament is a proton pump inhibitor (PPI).

4. The granules according to claim 3, wherein the PPI is a benzimidazole compound represented by the formula (I):



wherein ring A is an optionally substituted benzene ring,  $R^1$ ,  $R^2$  and  $R^3$  are the same or different, and represent a hydrogen atom, an optionally substituted alkyl group, an optionally substituted alkoxy group, or an optionally substituted amino group, and Y represents a nitrogen atom

or CH, or a salt thereof.

5. The granules according to claim 3, wherein the PPI is lansoprazole, omeprazole, rabeprazole, pantoprazole, leminoprazole, tenatoprazole (TU-199), or an optically  
5 active isomer thereof, or a pharmaceutically acceptable salt thereof.

6. The granules according to claim 3, wherein the PPI is lansoprazole, or an optically active isomer thereof, or a pharmaceutically acceptable salt thereof.

10 7. The granules according to claim 4, wherein the basic inorganic salt is contained in the principal ingredient layer in an amount of about 0.2 part by weight to about 0.6 part by weight based on 1 part by weight of the benzimidazole compound.

15 8. The granules according to claim 1 which are granules having a core, wherein the principal ingredient layer is formed on the core comprising at least one material selected from sucrose, starch, lactose and crystalline cellulose.

20 9. The granules according to claim 1, wherein the enteric coating layer contains an enteric water-soluble polymer.

10. The granules according to claim 9, wherein the enteric water-soluble polymer is a methacrylic acid  
25 copolymer.

11. The granules according to claim 1, wherein an average particle size of the granules is about 1000  $\mu\text{m}$  to about 2000  $\mu\text{m}$ .

12. The granules according to claim 4, wherein the  
5 benzimidazole compound is contained in an amount of about 12% by weight to about 40% by weight based on the total granules.

13. A granular preparation, a capsule, a tablet, an effervescent preparation or a suspension comprising the  
10 granules according to claim 1.

14. Granules comprising a principal ingredient layer containing a PPI in an amount of about 12% by weight to about 40% by weight based on the total granules, and one or more basic inorganic salts selected from the group  
15 consisting of basic salts of a sodium salt, a potassium salt, an aluminum salt, a magnesium salt and a calcium salt in an amount of about 0.2 part by weight to about 0.6 part by weight based on 1 part by weight of the PPI; an intermediate coating layer which is formed on the principal  
20 ingredient layer; and an enteric coating layer which is formed on the intermediate coating layer, and having an average particle size of about 1000  $\mu\text{m}$  to about 2000  $\mu\text{m}$ .

15. A granular preparation, a capsule or a tablet comprising the granules according to claim 14.

25 16. The preparation according to claim 15, which is a

preparation for treating or preventing peptic ulcer, Zollinger-Ellison syndromes, gastritis, reflux esophagitis, symptomatic gastroesophageal reflux disease (symptomatic GERD), NUD (non ulcer dyspepsia), stomach cancer, gastric MALT lymphoma, ulcer resulting from nonsteroidal antiinflammatory medicaments, or excess stomach acid or ulcer due to post-operation stress, a preparation for eradicating *Helicobacter pylori*, or a preparation for suppressing upper gastrointestinal hemorrhage due to peptic ulcer, acute stress ulcer, hemorrhagic gastritis or invasion stress.

17. Granules comprising a principal ingredient layer containing lansoprazole or an optically active isomer (R isomer) thereof in an amount of about 14% by weight to about 20% by weight based on the total granules, and one or more basic inorganic salts selected from the group consisting of basic salts of magnesium and calcium in an amount of about 0.2 part by weight to about 0.4 part by weight based on 1 part by weight of lansoprazole or an optically active isomer (R isomer) thereof; an intermediate coating layer which is formed on the principal ingredient layer; and an enteric coating layer which is formed on the intermediate coating layer, and having an average particle size of 1000  $\mu\text{m}$  to about 2000  $\mu\text{m}$ .

18. The granules according to claim 17, wherein the

basic inorganic salt is magnesium carbonate.

19. No. 3 to 5 capsules comprising 30 mg of lansoprazole per one capsule.

20. The capsule according to claim 19, wherein the granules according to claim 17 are filled.

21. No. 4 to 5 capsules comprising 15 mg of lansoprazole per one capsule.

22. The capsule according to claim 21, wherein the granules according to claim 17 are filled.

23. No. 1 to 3 capsules comprising 60 mg of lansoprazole or an optically active isomer (R isomer) thereof per one capsule.

24. The capsule according to claim 23, wherein the granule according to claim 17 is filled.

25. No. 2 to 4 capsules comprising 40 mg of an optically active isomer (R isomer) of lansoprazole per one capsule.

26. The capsule according to claim 25, wherein the granules according to claim 17 are filled.

27. No. 3 to 5 capsules comprising 30 mg of an optically active isomer (R isomer) of lansoprazole per capsule.

28. The capsule according to claim 27, wherein the granules as defined in claim 17 are filled.

29. A pharmaceutical composition which is a

combination of an antimicrobial agent and the granules  
according to claim 3.

Fetherstonhaugh  
Ottawa, Canada  
Patent Agents